

**Supplementary Table 3.** Seizure recurrence risk with sub lobar localization of calcification

Location	Seizure recurrence (n=35)	No recurrence (n=74)	<i>p</i> -value	Odds ratio (95% CI)
Frontal				
Orbitofrontal	0	0		
Polar	0	1 (1.4)	0.8	0.69 (0.027-17.6)
Dorsolateral	5 (14.3)	3 (4.1)	0.07	3.9 (0.88-17.5)
Ventrolateral	7 (20)	14 (18.9)	0.84	1.1 (0.4-3.06)
Superolateral	3 (8.5)	2 (2.7)	0.18	3.4 (0.54-21.48)
Mesial	1 (2.8)	14 (18.9)	0.05	0.12 (0.015-1.00)
Opercular	0	1 (1.4)	0.8	0.69 (0.027-17.6)
Parietal				
Anterolateral	2 (5.7)	6 (8.11)	0.65	0.68 (0.13-3.5)
Posterolateral	5 (14.27)	14 (18.91)	0.71	0.71 (0.25-2.17)
Mesial	6 (17.14)	6 (8.10)	0.15	2.42 (0.72-8.71)
Opercular	2 (5.7)	0	0.12	11.1 (0.51-238.02)
Temporal				
Anterolateral	0	1 (1.4)	0.8	0.69 (0.027-17.6)
Posterolateral	0	2 (2.7)	0.56	0.40 (0.019-8.73)
Mesial	1 (5.7)	0	0.26	6.47 (0.25-163.12)
Basal	0	0		
Opercular	0	0		
Occipital				
Anterolateral	1 (2.8)	0	0.25	6.4 (0.25-163.12)
Posterolateral	0	1 (1.4)	0.8	0.69 (0.027-17.6)
Mesial	1 (2.8)	9 (12.16)	0.14	0.21 (0.25-1.74)

Values are presented as number (%).